

APPENDIX 1
BRUXSCREEN-Q
Bruxism Screening Guide (BruxScreen)

Part I: Self-Report Questionnaire

Description:

This questionnaire evaluates the potential presence of bruxism and its related jaw symptoms. Bruxism is an activity of the jaw muscles that manifests in two distinct ways: as light or strong clenching of the jaw and teeth, or as grinding of the teeth. Clenching is a static activity, whereas during grinding, the lower jaw moves along the upper jaw. Grinding is often accompanied by grinding sounds. These activities can occur both during sleep and while awake. Generally, bruxism is a harmless activity; however, in some individuals, it may lead to potential consequences such as jaw pain, stiffness in the chewing muscles, or limitations in mouth opening.

You are kindly requested to complete the brief questionnaire below, enabling your dentist to gather more information regarding the potential presence of this jaw muscle activity. Once you have completed the questionnaire in full, please hand it to your dentist. Your dentist will subsequently conduct a brief examination of your jaw and teeth to confirm the presence of any conditions you have indicated in the questionnaire.

Thank you for your assistance!

Your Dentist

1. Bruxism

a. How often do you clench your teeth during sleep?

- ☐ Never
- ☐ Rarely
- ☐ Sometimes
- ☐ Often
- ☐ Always
- ☐ Don't know

b. How often do you grind your teeth during sleep?

- ☐ Never
- ☐ Rarely
- ☐ Sometimes
- ☐ Often
- ☐ Always
- ☐ Don't know

c. How often do you clench your teeth while awake?

- ☐ Never
- ☐ Rarely

- ☐ Sometimes
- ☐ Often
- ☐ Always
- ☐ Don't know

d. How often do you grind your teeth while awake?

- ☐ Never
- ☐ Rarely
- ☐ Sometimes
- ☐ Often
- ☐ Always
- ☐ Don't know

e. How often do you lightly press, touch, or bring your upper and lower teeth together, other than while eating?

- ☐ Never
- ☐ Rarely
- ☐ Sometimes
- ☐ Often
- ☐ Always
- ☐ Don't know

f. How often do you tightly contract or tense your jaw muscles without clenching or bringing your teeth together while awake?

- ☐ Never
- ☐ Rarely
- ☐ Sometimes
- ☐ Often
- ☐ Always
- ☐ Don't know

2. Jaw Symptoms

a. How often do you experience pain, discomfort, tenderness, fatigue, tension, or stiffness in your temple, face, jaw, or jaw joint areas?

Symptom	Upon Waking Up	At Any Other Time During the Day
Pain	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Discomfort	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Tenderness	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Fatigue	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Tension	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often

Symptom	Upon Waking Up	At Any Other Time During the Day
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Stiffness	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know

b. How often do you experience pain, discomfort, tenderness, fatigue, tension, or stiffness in your temple, face, jaw, or jaw joint areas when opening your mouth or during chewing?

Symptom	Upon Waking Up	At Any Other Time During the Day
Pain	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Discomfort	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Tenderness	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often

Symptom	Upon Waking Up	At Any Other Time During the Day
Fatigue	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
Tension	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know
Stiffness	☐ Never	☐ Never
	☐ Rarely	☐ Rarely
	☐ Sometimes	☐ Sometimes
	☐ Often	☐ Often
	☐ Always	☐ Always
	☐ Don't know	☐ Don't know

c. How often do you experience jaw locking or sticking?

(i) During meals:

- ☐ Never
- ☐ Rarely
- ☐ Sometimes
- ☐ Often

- ☐ Always
- ☐ Don't know

(ii) At other times:

- ☐ Never
- ☐ Rarely
- ☐ Sometimes
- ☐ Often
- ☐ Always
- ☐ Don't know

BRUXSCREEN-C

Bruxism Screening Guide (BruxScreen)

Part II: Clinical Assessment Form

Explanation

This clinical assessment tool aims to identify signs that may be associated with bruxism. First, the facial contours are examined for potential hypertrophy of the masseter muscles. This examination is performed both when the muscles are relaxed and when they are contracted. In case of uncertainty, select "absent." Second, the oral cavity is examined for signs that may be associated with bruxism. Similarly, in case of doubt, select "absent." Third, the teeth are evaluated for tooth wear. Please follow the scoring rules provided. If uncertain between two scores, choose the lower one. Additionally, specify whether the observed tooth wear is primarily mechanical (e.g., due to tooth-to-tooth contact), chemical, or a combination of both.

1. Extraoral Examination

a. Masseter muscle hypertrophy (when muscles are relaxed):

☐ Absent

☐ Present

b. Masseter muscle hypertrophy (when muscles are contracted):

☐ Absent

☐ Present

2. Intraoral Examination (Non-Dental Tissues)

a. Lip (indentations caused by teeth):

☐ Absent

☐ Present

b. Cheek (linea alba - buccal mucosa ridging):

☐ Absent

☐ Present

c. Tongue (scalloped tongue - indentations caused by teeth):

☐ Absent

☐ Present

d. Tongue (traumatic lesions):

☐ Absent

☐ Present

e. Alveolar bone (exostoses/tori):

☐ Absent

☐ Present

3. Intraoral Examination of Teeth

a. Occlusal/Incisal Wear Per Sextant:

- **(0):** No visible wear
- **(1):** Wear visible within enamel boundaries
- **(2):** Exposure of dentin with a loss of $\leq 1/3$ of the clinical crown height
- **(3):** Loss of $> 1/3$ but $< 2/3$ of the clinical crown height
- **(4):** Loss of $\geq 2/3$ of the clinical crown height

b. Palatal Wear in Sextant 2:

- **(0):** No visible wear
- **(1):** Wear limited to enamel
- **(2):** Wear reaching dentin boundaries

Wear Assessment Per Sextant:

Sextant	0	1	2	3	4
Sextant 1 – Occlusal	☐	☐	☐	☐	☐
Sextant 2 – Incisal	☐	☐	☐		
Sextant 3 – Occlusal	☐	☐	☐	☐	☐
Sextant 2 – Palatal	☐	☐	☐		
Sextant 6 – Occlusal	☐	☐	☐	☐	☐
Sextant 5 – Incisal	☐	☐	☐		
Sextant 4 – Occlusal	☐	☐	☐	☐	☐

c. Cause of Tooth Wear:

☐ Primarily mechanical

☐ Primarily chemical

☐ Both mechanical and chemical